



VACCINE SCHEDULES

Children in the United States are recommended to be vaccinated against 15 diseases (those recommended for all children). Additional vaccines are added based on shared decision making or specific conditions, and in the National Schedules of each country based on their prevalence or endemic infections. For example, the BCG vaccine is part of the routine immunization schedule of Saudi Arabia and the United Arab Emirates.

This guide will go over **the vaccines as recommended by American Academy of Pediatrics**. In the next pages, you will see an immunizations schedule from birth through 18 years of age. You can print this and take it to your pediatrician visits for reference, or if you have any questions. Following the schedule, you will find a notes section covering what each vaccine protects against, and some special conditions depending on the vaccine.

Please note that the included schedule is intended to cover **routine vaccinations for healthy children**. There are certain vaccines that are recommended for special populations that are not included in this guide, kindly confirm with your physician. These have been excluded for simplicity purposes, and they can be shared upon request to pedsandco@gmail.com.

Saudi Arabia National
Immunization Schedule



UAE National
Immunization
Schedule



Vaccine	Birth	1 mo	2 mo	4 mo	6 mo	8 mo	9 mo	12 mo	15 mo	18 mo	19–23 mo	2–3 yr	4–6 yr	7–10 yr	11–12 yr	13–15 yr	16 yr	17–18 yr
RSV protection (antibody shot)	1 dose before 8 months of age, entering their first RSV season																	
Hepatitis B (HepB)	1st	2nd			3rd													
Rotavirus (RV)			1st	2nd	3rd †													
Diphtheria, tetanus, pertussis (DTaP)			1st	2nd	3rd				4th				5th					
Hib			1st	2nd	3rd †			3rd or 4th										
Pneumococcal (PCV)			1st	2nd	3rd			4th										
Polio (IPV)			1st	2nd	3rd								4th					
COVID-19					Routine: 1 or 2 doses													
Influenza (flu)					Every year, from 6 months of age													
Measles, mumps, rubella (MMR)								1st					2nd					
Chickenpox (Varicella)								1st					2nd					
Hepatitis A (HepA)								2 doses, ≥6 months apart										
Tdap															1 dose			
HPV														2-dose series: start at 9–12 yr				
Meningococcal ACWY															1st			
Meningococcal B																	2nd	2-dose series: shared decision

Recommended for all healthy children

Shared decision with your doctor

† Rotavirus and Hib: whether a 3rd dose is given at 6 months depends on which brand is used. Both schedules are complete. **Ages** are read as “through”: 12–15 mo means 12 months through 15 months.

Peds&Co · @pedsandco · Free extract from ?Vaccines Must Know!?